

NORDIMET® PRE-FILLED SYRINGE Brief injection instructions

Please note

- Injections take place just once a week. On a specific day.
- Only use the syringe before the expiration date and if the solution is yellow and clear.
- Potential air bubbles in the solution are not harmful and are not removed before injection.
- These instructions do not supersede the patient information leaflet. Please read it carefully.

Preparing the injection





- Clean your hands thoroughly.
- Lay out the syringe and alcohol pads ready.
- Remove the syringe from the packaging by the cylinder, do not touch the plunger or the needle cap.
- Select an injection site (lower stomach, a few centimetres away from the belly button, or on the front of the thigh).
- Clean the injection site with the alcohol pads included.
- Consider changing the injection site for each injection.
- Tip: Use an injection calendar to document the date and injection site of the weekly injection.

Performing the injection









- Hold the syringe by the cylinder and pull the cap right off.
- Do not touch the plunger yet.
- ▶ Do not touch the needle before the injection.
- ▶ Hold the syringe in your hand like a pen.
- ▶ Form a skin fold at the selected injection site with the thumb and index finger of your other hand.
- ▶ Insert the needle fully into the skin fold at a 90-degree angle.
- Push the plunger down all the way with your thumb until the syringe is empty.
- Take your thumb away from the plunger and remove the syringe from the skin fold.
- ▶ The safety system automatically retracts the needle.
- ▶ Release the skin fold.



NORDI